REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and

			ense, Executive Service Directors collection of information if it does [HE ABOVE ORGANIZA]		spondents sho y valid OM8 c	duld be, aware that notwithstanding any other provision of law, no control number:
1. REPORT D	ATE (DD-MM-YY 1/26/2017		ORT TYPE Poster/Abs		3. DATES COVERED (From - To) 10/26/2017-10/30/2017	
4. TITLE AND Multiple co-me		n patient with N	Mast Cell Activation Synd	rome	5a. CO	NTRACT NUMBER
					5b. GR	ANT NUMBER
					5c. PRO	OGRAM ELEMENT NUMBER
6. AUTHOR(S	· · · · · · · · · · · · · · · · · · ·				5d. PRO	OJECT NUMBER
Maj Sofia Szar						
					5e. 1AS	SK NUMBER
					5f. WO	RK.UNIT NUMBER
59th Clinical R	NG ORGANIZAT lesearch Division Hall Loop, Bldg	- *	ND ADDRESS(ES)			8. PERFORMING ORGANIZATION REPORT NUMBER
JBSA-Lackland, TX 78236-9908 210-292-7141						1.7341
59th Clinical R 1100 Willford	escarch Division Hall Loop, Bidg	4430	//E(S) AND ADDRESS(ES	š) .		10. SPONSOR/MONITOR'S ACRONYM(S)
JBSA-Lacklan 210-292-7141	d, TX 78236-990	8				11. SPONSOR/MONITOR'S REPORT NUMBER(S)
	ION/AVAILABIL Jublic release, Di					
13. SUPPLEME	NTARY NOTES					
Sofia M. Szari Wilford Hall A Disclaimer: Th	orbid conditions MD, and James mbulatory Surgi	M. Quinn. MD cal Center, Joint crtions herein a		kland AFB, TX		e construed as reflecting the views of the
intolerance and	Vor postural orth	ostatic tachycard	lia syndrome (POTS) in l	imited reports.	A single	Tissue Disorders (CTD) and orthostatic report has also identified kindreds of patients with yehiatric findings, atopy, and gastrointestinal diso
15. SUBJECT	ERMS	· · · · · · · ·				
16. SECURITY CLASSIFICATION OF: a. REPORT b. ABSTRACT c. THIS PAGE			17. LIMITATION OF ABSTRACT	18. NUMBER OF		ME OF RESPONSIBLE PERSON Longoria
21 1.20 A131	<u></u>	o. mo i Ade	ŬU	PAGES		EPHONE NUMBER (Include area code)

Multiple co-morbid conditions in patient with Mast Cell Activation Syndrome


Maj Sofia M. Szari, MD and James M. Quinn, MD Wilford Hall Ambulatory Surgical Center, Joint Base San Antonio - Lackland AFB, TX*


Introduction

Mast cell activation disorders (MCAD) have been associated with Connective Tissue Disorders (CTD) and orthostatic intolerance and/or postural orthostatic tachycardia syndrome (POTS) in limited reports. A single report has also identified kindreds of patients with an autosomal dominant pattern of increased serum tryptase and variably associated CTD, neuropsychiatric findings, atopy, and gastrointestinal disorders. This abstract describes an unusual case of MCAD; Ehlers-Danlos Hypermobility Syndrome (EDHS); orthostatic intolerance; anxiety/ADHD; intermittent diarrhea; and a carcinoid tumor. This cluster of co-morbid conditions with MCAD has not been described to occur in a single patient.

Case Report

A 24 year-old female with a history of fire ant hypersensitivity on immunotherapy, anxiety/ADHD, episodes of orthostatic intolerance, and intermittent diarrhea was found to have an elevated baseline tryptase of 13.7 mcg/L persisting at 13.8 mcg/L on repeat. She was evaluated by hematology/oncology and diagnosed with a MCAD with a low likelihood for systemic disease and offered a bone marrow biopsy. Dermatology consultation concurred with MCAD without evidence of systemic or cutaneous disease. During her evaluation for MCAD, she was diagnosed with EDHS after formal evaluation by geneticist. Shortly after, she underwent appendectomy for acute abdominal pain. Her surgical pathology of the removed appendix revealed a grade 1 carcinoid tumor with negative margins, negative lymph nodes and no inflammation (net G1, path pTaNO).





Discussion

This case illustrates the many potential conditions that can be associated with mast cell disorders and CTD. Connective tissue, namely the components of the extracellular matrix, is known to have affects on mast cells including their migration. It stands to reason that there could be associations with CTD, including orthostatic intolerance and Ehlers-Danlos Syndrome with mast cell disorders. There has been a proposed disease cluster of EDS, POTS and MCAD, which our patient would qualify. She additionally has 2 other associated findings of neuropsychiatric condition (anxiety/ADHD) and gastrointestinal symptoms (intermittent idiopathic diarrhea). Her recent excision of a carcinoid tumor has not been described to date in association with MCAD and is potentially a newly linked association vs incidental.

Conclusion

This case emphasizes the need to evaluate for co-morbid conditions in patients with MCAD. The several conditions associated with mast cell disorders in this patient may represent a newly identified cluster.

References

- Seneviratne SL, Maitland A, Afrin L. 2017. Mast cell disorders in Ehlers-Danlos syndrome. Am J Med Genet Part C Semin Med Genet 175C:228-236.
- Cheung I, Vadas P. 2015. A new disease cluster: Mast cell activation syndrome, postural orthostatic tachycardia syndrome, and Ehlers-Danlos syndrome. J Allergy Clinical Immunol 135(2):AB65.
- Louisias M, Silverman S, Maitland AL. 2013. Prevalence of allergic disorders/mast cell activation syndrome in patients with Ethiers-Danios syndrome. Annais of allergy, asthma & immunol, A12. Baltimore, MD, USA: American College of Allergy, Asthma & immunology.
- Fikree A, Aktar R, Grahame R, Hakim AJ, Morris JK, Knowles CH, Aziz Q. 2015. Functional gastrointestinal disorders are associated with the joint hypermobility syndrome in secondary care: A case-control study. Neurogastroenterol Molil 27:5696-579.
- Sinibaldi L, Ursini G, Castori M. 2015. Psychopathological manifestations of joint hypermobility and joint hypermobility syndrome/Ehlers-Danlos syndrome, hypermobility type: The link between connective tissue and psychological distress revised. Am J Med Genet Part C Semin Med Genet 196C-97-108.
- Garland EM, Celedonio JE, Raj SR. 2015. Postural tachycardia syndrome: Beyond orthostatic intolerance. Curr Neurol Neurosci Rep 15:60.

*The opinions or assertions herein are the private views of the authors and are not to be construed as reflecting the views of the Department of the Air Force or the Department of Defense.

Multiple co-morbid conditions in patient with Mast Cell Activation Syndrome

Sofia M. Szari, MD, and James M. Quinn, MD

Wilford Hall Ambulatory Surgical Center, Joint Base San Antonio – Lackland AFB, TX

Disclaimer: The opinions or assertions herein are the private views of the authors and are not to be construed as reflecting the views of the Department of the Air Force or the Department of Defense.

INTRODUCTION: Mast cell activation disorders (MCAD) have been associated with Connective Tissue Disorders (CTD) and orthostatic intolerance and/or postural orthostatic tachycardia syndrome (POTS) in limited reports. A single report has also identified kindreds of patients with an autosomal dominant pattern of increased serum tryptase and variably associated CTD, neuropsychiatric findings, atopy, and gastrointestinal disorders. This abstract describes an unusual case of MCAD; Ehlers-Danlos Hypermobility Syndrome (EDHS); orthostatic intolerance; anxiety/ADHD; intermittent diarrhea; and a carcinoid tumor. This cluster of co-morbid conditions with MCAD has not been described to occur in a single patient:

CASE PRESENTATION: A 24 year-old female with a history of fire ant hypersensitivity on immunotherapy, anxiety/ADHD, episodes of orthostatic intolerance, and intermittent diarrhea was found to have an elevated baseline tryptase of 13.7 mcg/L persisting at 13.8 mcg/L on repeat. She was evaluated by hematology/oncology and diagnosed with a MCAD with a low likelihood for systemic disease and offered a bone marrow biopsy. Dermatology consultation concurred with MCAD without evidence of systemic or cutaneous disease. During her evaluation for MCAD, she was diagnosed with EDHS. Shortly after, she underwent appendectomy for acute abdominal pain. Her surgical pathology revealed a grade 1 carcinoid tumor with negative margins, negative lymph nodes and no inflammation (net G1, path pTaNO).

DISCUSSION: This case emphasizes the need to evaluate for co-morbid conditions in patients with MCAD. The several conditions associated with mast cell disorders in this patient may represent a newly identified cluster.

References

- Seneviratne SL, Maitland A, Afrin L. 2017. Mast cell disorders in Ehlers—Danlos syndrome. Am J Med Genet Part C Semin Med Genet 175C:226—236.
- 2. Cheung I, Vadas P. 2015. A new disease cluster: Mast cell activation syndrome, postural orthostatic tachycardia syndrome, and Ehlers-Danios syndrome. J Allergy Clinical Immunol 135(2):AB65.
 - Louisias M, Silverman S, Maitland AL. 2013. Prevalence of allergic disorders/mast cell activation syndrome in patients with Ehlers-Danlos syndrome. Annals of allergy, asthma & immunol, A12. Baltimore, MD, USA: American College of Allergy, Asthma & Immunology.
- 3. Fikree A, Aktar R, Grahame R, Hakim AJ, Morris JK, Knowles CH, Aziz Q. 2015. Functional gastrointestinal disorders are associated with the joint hypermobility syndrome in secondary care: A case-control study. Neurogastroenterol Motil 27:569–579.
- 4. Sinibaldi L, Ursini G, Castori M. 2015. Psychopathological manifestations of joint hypermobility and joint hypermobility syndrome/Ehlers—Danlos syndrome, hypermobility type: The link between connective tissue and psychological distress revised. Am J Med Genet Part C Semin Med Genet 169C:97—106.

4	
intolerance. Curr Meurol Meurosci Rep 15:60.	
Garland EM, Celedonio JE, Raj SR. 2015. Postural tachycardia syndrome: Beyond orthostatic	٦.